

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: A61K 47/26, 38/24	A1	(11) International Publication Number: WO 96/29095 (43) International Publication Date: 26 September 1996 (26.09.96)
(21) International Application Number: PCT/EP95/01055 (22) International Filing Date: 21 March 1995 (21.03.95) (71) Applicant (for all designated States except US): APPLIED RESEARCH SYSTEMS ARS HOLDING N.V. [NL/NL]; John B. Gonsiraweg 14, Curaçao (AN). (72) Inventors; and (75) Inventors/Applicants (for US only): SAMARITANI, Fabrizio [IT/IT]; Via Luigi Chiala, 130, I-00139 Rome (IT). NATALE, Patrizia [IT/IT]; Via Tirso, 80, I-00198 Rome (IT). (74) Agent: VANNINI, Mario; Istituto Farmacologico Serrone S.p.A., Via Casilina, 125, I-00176 Rome (IT).		(81) Designated States: AU, BR, BY, CA, CN, FI, JP, KR, KZ, MX, NO, RU, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: HCG LIQUID FORMULATIONS (57) Abstract The invention refers to liquid pharmaceutical compositions containing hCG stabilised with a polyalcohol or a non-reducing sugar. Preferably, the compositions are stabilised with mannitol. In the preferred embodiments such compositions are aqueous solutions in a phosphate buffer at pH 7. Such compositions are ready to be injected and, therefore, the step of reconstitution of the lyophilised powder is avoided, thus simplifying the way of use.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

HCG LIQUID FORMULATIONS

The present invention relates to gonadotropin containing liquid pharmaceutical compositions. More precisely, it concerns liquid formulations of hCG (human Chorionic Gonadotropin) stabilised with a polyalcohol or a non-reducing sugar.

It is known that highly purified proteins easily undergo degradation, even due to the contact with atmospheric agents. This characteristic is even more evident for proteins produced by recombinant DNA techniques.

Such proteins are usually stabilised with saccharides, such as lactose, or with mannitol, or else with proteins or aminoacids, such as albumin and glycin.

The injectable stabilised formulations of gonadotropins are obtained with a process which includes always a step of lyophilisation to obtain a dry powder; in such a way the stabilised formulations are able to maintain a longer cycle life, even if stored at room temperature.

WO 93/11788 describes lyophilised gonadotropin-containing pharmaceutical compositions stabilised with sucrose, alone or in combination with other stabilising agents. In said patent application it is shown that the stability provided to the lyophilised compositions under study by sucrose was better than that provided by lactose or mannitol.

No liquid stabilised formulations of gonadotropins have been described until now. It is highly desirable to obtain such liquid formulations so as to have the compositions ready to be injected and to avoid the reconstitution of the lyophilised powder, thus simplifying the way of use.

We have surprisingly found that it is possible to obtain such liquid stabilised formulations.

The main object of the present invention is to provide a liquid pharmaceutical composition containing hCG stabilised with a polyalcohol or a non-reducing sugar. Preferably the polyalcohol is mannitol and the non-reducing sugar is sucrose. More preferably the liquid formulations of the invention are stabilised with mannitol.

The solution is preferably a buffered aqueous solution and the buffer according to the invention is selected from the group consisting of

phosphate, acetate or succinate buffer. The preferred buffer is phosphate and the pH is preferably 7.00.

The hCG is preferably recombinant and can be prepared, for example, by expression in CHO (Chinese Hamster Ovary) cells, transformed
5 with the corresponding DNA, according to the technique described in European Patent 160699.

A further object of the present invention is to provide a process for the preparation of said liquid pharmaceutical composition comprising diluting a hCG bulk solution in a buffer solution containing the excipients.

10 Still another object of the present invention is to provide a form of presentation of said liquid pharmaceutical composition comprising such formulation hermetically closed in a sterile condition in a container suitable for the storage before the use.

In order to optimise the stability of the hCG formulations of the invention a series of preliminary experiments have been carried out with
15 different buffers at various pH, ionic strength, dielectric constant and concentration of rec-hCG.

In order to evaluate the effect of pH and of the buffer, 0.01 M solutions of phosphate, succinate or acetate buffers were prepared with
20 water for injection. The pH was adjusted to 6.0, 7.0 or 8.0 with NaOH 1 M. The bulk solution of rec-hCG was added to the buffer systems to obtain solutions at 5,000 IU/ml. The solutions were then filtered and poured into 3 ml glass vials. The composition of the formulations thus prepared is reported in Table 1. The accelerated stability of these formulations has been studied,
25 so that the stability of the same can be foreseen when they are stored in containers at room temperature, through the extrapolation of the data at higher temperatures. In this case the samples were stored at 40° and 50°C and the stability of rec-hCG was checked by determining its purity by HPSEC analyses according to the following standard conditions:

30 Phase A	0.1 M phosphate pH 6.7 + 0.1 M Na ₂ SO ₄
Isocratic conditions	100% phase A
Column	TSK G 2000 SWXL
Flow Rate	0.5 ml/min
UV Detector	214 nm
35 Injection Volume	20 µl (10,000 IU strength)

40 µl (5,000 IU strength)

Table 3 reports the percentage of rec-hCG monomer peak determined by HPSEC. The results show that the solutions at pH 6.0 and 8.0 are less stable in comparison with the solutions at pH 7.0 and that no remarkable stability differences were observed among the buffers.

The effect of the ionic strength was evaluated on rec-hCG 5,000 IU/ml solutions, prepared with phosphate and succinate buffers 0.01M at pH 7.0, adjusted with NaCl to the following values of osmolality: 150, 300 and 400 mOsm. The composition of the formulations is reported in Table 2. The samples were stored at 4°, 25°, 40° and 50°C and tested for the stability of rec-hCG by HPSEC. The results, reported in table 4, show that the increase of ionic strength negatively affects the stability of rec-hCG.

The effect of the dielectric constant was evaluated on 5,000 IU/ml solutions of rec-hCG, prepared with phosphate and succinate buffers 0.01 M at pH 7, containing 5, 10 and 15% propylene glycol. The composition of the formulations is reported on Table 2. The samples were stored at 4°, 25°, 40° and 50°C and tested for the stability of rec-hCG by HPSEC. The results, reported in Table 4, show that increasing the percentage of propylene glycol negatively affects the stability of rec-hCG.

In order to evaluate the effect of the rec-hCG concentration, the stability at 50°C of the solutions in phosphate buffer 0.01 M at pH 7.0, containing respectively 2,500, 5,000, 7,500 and 10,000 IU/ml of rec-hCG was monitored by HPSEC for 2 weeks. The results reported in Table 5 showed that the stability was higher for the more concentrated solutions.

In order to compare the effects of various stabilisers and/or excipients on the stability of rec-hCG, six liquid formulations, in phosphate buffer 0.01 M at pH 7.0 containing 10,000 IU/ml rec-hCG were prepared, as a first step. Sucrose, glycine, glucose, mannitol, lactose and NaCl were used, as stabilisers/excipient. The composition of the formulations is reported in Table 6. These formulations were submitted to the stability tests by storing samples at 4°, 25°, 40° and 50°C and tested by a Bioassay and HPSEC. Subsequently, based on the results of said first step, four lots of two selected liquid formulations were prepared, using as stabilisers sucrose and mannitol. Table 7 reports the composition of such formulations.

The Bioassay has been carried out in accordance with the European Pharmacopocia Monograph.

In Table 8 the HPSEC stability data are reported and in Table 9 the values of bioactivity are reported. The results showed the following:

- 5 1. the bioactivity of the formulations containing glucose and lactose remarkably decreased at 50°C after 1 week storage. Also monomer peak was lower compared to that measured in the other formulations.
2. in the presence of glycine and NaCl a more evident decrease of bioactivity and of purity was measured in comparison to the formulations containing sucrose and mannitol. Also in this case the decrease of the percentage of the rec-hCG monomer peak, was not due to the formation of aggregates forms, but to the increase of free subunits.

Tables 10 and 11 report the purity determined by HPSEC for the 5,000 and 10,000 IU strength respectively. Data show that even after three weeks at 50°C the purity is higher in the formulations containing mannitol compared to the formulations containing sucrose. Tables 12 and 13 report the purity of the α subunit determined by reverse phase HPLC after 1 week storage at 50°C for the sucrose and mannitol formulations. The data confirm the better stability of the formulation containing mannitol in comparison to that containing sucrose.

Reverse Phase HPLC analyses have been performed with the following standard conditions:

Phase A	1 ml TFA in 1 liter of bidistilled water		
Phase B	0.79 ml TFA in 1 liter of acetonitrile		
25 Gradient conditions	time	A%	B%
	0	85	15
	20'	60	40
	21'	20	80
	22'	85	15
30 Column	Aquapore RP 300 25 cm		
Column temperature	40°C		
Flow Rate	1 ml/min		
UV detector	214 nm		
Injection volume	10 μ l		

In the Tables 14 and 15 the results of the bioactivity assay are reported. No appreciable bioactivity decrease was observed after 24 weeks at 4° and 25°C in the mannitol formulation.

According to the present invention, the liquid pharmaceutical compositions contain from 1,000 to 40,000 IU/ml, preferably 10,000 IU/ml, of hCG and from 10 to 180 mg/l, preferably 54.6 mg/l, of mannitol in a 0.01 M buffer solution.

EXAMPLES OF PHARMACEUTICAL MANUFACTURING

Materials: Phosphoric acid 85% RPE ACS (Carlo Erba); Mannitol DAB, Ph Eur BP, FU, USP, FCC, E421 (Merck), NaOH 1 M (Merck), water for injections.

The primary container for the formulated vials consists of: 3 ml glass vials (DIN 2R) (borosilicate glass type I), Rubber closures (Pharmagummi W1816 V50), Aluminium rings and flip off caps (Pharma Metal GmbH).

Preparation of rec-hCG solution containing mannitol

The phosphoric acid (0.98 g) is added to the water for injections (600 ml). If necessary, the pH is adjusted to 7.0 with NaOH 1 M. Mannitol (54.6 g) is added to the phosphoric acid solution and the pH is again checked and, if necessary, adjusted to 7.00 ± 0.2 with NaOH 1 M or with phosphoric acid diluted 1:5. The rec-hCG bulk (10 MIU or 20 MIU, if the final desired strenght is 5,000 or 10,000 IU respectively) is then added to the excipient solution and the pH is again checked and, if necessary, adjusted to 7.00 ± 0.2 with NaOH 1 M or with phosphoric acid diluted 1:5.

The solution is brought to 1 liter with water for injections. Such solution is then filtered through a 0.22 μm Millipak 20 filter under a pressure not higher than 1.5 atm, under laminar flow, collecting the solution into a flask and stirring gently for about 1 minute.

The vials are then filled up with 0.5 ml of the rec-hCG solution.

TABLE 1 - COMPOSITION OF r-hCG SOLUTIONS**pH/buffer effect**

Acetate buffer solution	Amount/ml
r-hCG bulk	5000 IU
<u>Acetic acid glacial</u>	0.6 mg
NaOH 1M	q.s. to pH 6.0,7.0,8.0
Succinate buffer solution	
r-hCG bulk	5000 IU
<u>Succinic acid</u>	1.18 mg
NaOH 1M	q.s. to pH 6.0,7.0,8.0
Phosphate buffer solution	
r-hCG bulk	5000 IU
<u>Phosphoric acid 85%</u>	0.98 mg
NaOH 1M	q.s. to pH 6.0,7.0,8.0

Filling volume: 1 ml

TABLE 2 - COMPOSITION OF r-hCG SOLUTIONS

Ionic strength/dielectric constant

Lot	r-hCG	NaCl	Prop. glyc.	Phosp. buffer 0.01 pH 7.0	Succinate buffer 0.01 M pH 7.0
Fos/7.0/PG 5	5000 IU/ml	-	50 mg/ml	q.s. to 1 ml	-
Fos/7.0/PG 10	5000 IU/ml	-	100 mg/ml	q.s. to 1 ml	-
Fos/7.0/PG 15	5000 IU/ml	-	150 mg/ml	q.s. to 1 ml	-
Suc/7.0/PG 5	5000 IU/ml	-	50 mg/ml	-	q.s. to 1 ml
Suc/7.0/PG 10	5000 IU/ml	-	100 mg/ml	-	q.s. to 1 ml
Suc/7.0/PG 15	5000 IU/ml	-	150 mg/ml	-	q.s. to 1 ml

Filling volume: 1 ml

FOS = Phosphate buffer

SUC = Succinate buffer

7.0 = pH 7.0

PG 5 = propylene glycol 5%

PG 10 = propylene glycol 10%

PG 15 = propylene glycol 15%

TABLE 2 (CONT.)

LOT	r-hCG	NaCl	Prop. glyc.	Phosp. buffer 0.01 pH 7.0	Succinate buffer 0.01 M pH 7.0
Fos/7.0/150	5000 IU/ml	4.4 mg/ml	-	q.s. to 1 ml	-
Fos/7.0/300	5000 IU/ml	8.8 mg/ml	-	q.s. to 1 ml	-
Fos/7.0/400	5000 IU/ml	11.7 mg/ml	-	q.s. to 1 ml	-
Suc/7.0/150	5000 IU/ml	4.4 mg/ml	-	-	q.s. to 1 ml
Suc/7.0/300	5000 IU/ml	8.8 mg/ml	-	-	q.s. to 1 ml
Suc/7.0/400	5000 IU/ml	11.7 mg/ml	-	-	q.s. to 1 ml

Filling volume: 1 ml

150, 300, 400: osmolarity

FOS = Phosphate buffer

SUC = Succinate buffer

7.0 = pH 7.0

TABLE 3 - r-hCG PURITY (%)

HPSEC DATA

pH/Buffer effect

LOT	T=0	50 °C			40° C	
		1 W	3 W	5 W	3 W	5 W
ACE/6	100	95.85	92.70	84.99	97.51	94.10
ACE/7	100	96.62	93.26	88.02	97.27	94.05
ACE/8	100	96.51	92.70	87.10	97.45	95.12
SUC/6	100	94.56	91.28	82.11	96.92	93.11
SUC/7	100	95.78	94.20	88.05	96.91	93.99
SUC/8	100	95.36	90.12	83.00	97.61	94.02
FOS/6	100	94.10	90.76	81.00	97.50	93.00
FOS/7	100	96.09	93.12	86.93	96.72	93.74
FOS/8	100	94.21	82.52	74.96	96.77	93.55

W = week

ACE = acetate buffer

SUC = succinate buffer

FOS = phosphate buffer

6/7/8 = pH 6.0, 7.0, 8.0

TABLE 4 - r-hCG PURITY (%)
HPSEC DATA

Ionic strength/dielectric constant		50°C				40°C		25°C	4°C
LOT									
	T=0	1 W	2 W	4 W		3 W	6 W	6 W	4 W
Fos/7.0/PG 5	100	91.9	85.7	80.8		96.5	94.3	100	100
Fos/7.0/PG 10	100	91.9	81.0	77.7		93.9	93.9	100	100
Fos/7.0/PG 15	100	89.2	79.3	76.2		94.4	93.8	100	100
Suc/7.0/PG 5	100	90.6	84.3	-		91.7	-	-	-
Suc/7.0/PG 10	100	88.9	81.4	-		94.1	-	-	-
Suc/7.0/PG15	100	89.3	79.9	-		93.5	-	-	-

- = not tested

FOS = Phosphate buffer

SUC = Succinate buffer

7.0 = pH 7.0

PG 5 = propylene glycol 5%

PG 10 = propylene glycol 10%

PG 15 = propylene glycol 15%

W= week

TABLE 4 (CONT.)

LOT	T=0	50°C				40°C			25°C	4°C
		1 W	2 W	4 W	6 W	3 W	6 W	4 W	6 W	4 W
Fos/7.0/150	100	88.5	79.2	72.2	93.0	93.0	93.0	100	100	100
Fos/7.0/300	100	80.5	75.0	67.9	93.4	93.4	92.1	100	100	100
Fos/7.0/400	100	81.5	74.8	67.4	94.6	94.6	93.8	100	100	100
Suc/7.0/150	100	83.1	87.4	-	94.3	-	-	-	-	-
Suc/7.0/300	100	82.4	76.7	-	93.9	-	-	-	-	-
Suc/7.0/400	100	81.8	74.6	-	93.5	-	-	-	-	-

- = not tested

FOS = Phosphate buffer

SUC = Succinate buffer

7.0 = pH 7.0

150, 300, 400 = osmolality

W = week

TABLE 5 - r-hCG PURITY (%)**HPSEC DATA**

concentration effect

LOT		50°C		
		T=0	1 W	2 W
Fos /2500		100	87.3	84.0
Fos/5000		100	90.8	89.1
Fos/7500		100	92.9	89.8
Fos/10000		100	92.5	90.9

Fos/2500: 2,500 IU/ ml of r-hCG
Fos/5000 :5,000 IU/ ml of r-hCG
Fos/7500: 7,500 IU/ ml of r-hCG
Fos/10000:10,000 IU/ml of r-hCG

TABLE 6 - LIQUID FORMULATIONS

Vial composition

COMPONENTS/ LOT	r-hCG/ SAC	r-hCG/ MAN	r-hCG/ GLY	r-hCG/ GLU	r-hCG/ LAT	r-hCG/ NaCl
r-hCG IU/ml	10,000	10,000	10,000	10,000	10,000	10,000
SUCROSE mg/ml	102.6	-	-	-	-	-
MANNITOL mg/ml	-	54.6	-	-	-	-
GLYCINE mg/ml	-	-	22.52	-	-	-
GLUCOSE mg/ml	-	-	-	54.6	-	-
LACTOSE mg/ml	-	-	-	-	102.6	-
NaCl mg/ml	-	-	-	-	-	9.0

Buffer : H₃PO₄ 0.01 M , pH 7.0
 Filling volume: 0.5 ml

TABLE 7 - LIQUID FORMULATIONS**Vial composition**

COMPONENT	UNIT	r-hCG/5000/S01	r-hCG/10000/S01
r-hCG	IU/ml	10,000	20,000
SUCROSE	mg/ml	102.6	102.6
O. PHOSPHORIC ACID	mg/ml	0.98	0.98
SODIUM HYDROXIDE		q.s. to pH 7.0	q.s. to pH 7.0

COMPONENT	UNIT	r-hCG/5000/M01	r-hCG/10000/M01
r-hCG	IU/ml	10,000	20,000
MANNITOL	mg/ml	54.6	54.6
O. PHOSPHORIC ACID	mg/ml	0.98	0.98
SODIUM HYDROXIDE		q.s. to pH 7.0	q.s. to pH 7.0

Filling volume : 0.5 ml

TABLE 8 - COMPATIBILITY WITH DIFFERENT EXCIPIENTS

HPSEC stability data: purity (%)

LOT	30°C						40°C						25°C						4°C		
	T=0	1 W	2 W	6 W	2 W	4 W	6 W	11 W	4 W	6 W	8 W	11 W	6 W	8 W	11 W	8 W	11 W	8 W	12 W		
FOS/SAC	100	94.1	90.3	83.0	98.0	95.5	96.1	94.8	100	100	100	100	100	100	100	100	100	100	100		
FOS/GLY	100	94.2	90.4	81.5	97.5	96.3	95.5	95.5	100	100	100	100	100	100	100	100	100	100	100		
FOS/GLU	100	85.0	74.9	N.T	88.0	N.T	N.T	N.T	N.T	N.T	N.T	N.T	N.T	N.T	N.T	N.T	N.T	N.T	N.T		
FOS/MAN	100	94.0	91.7	83.5	97.9	97.1	95.8	95.4	N.T	100	100	100	100	100	100	100	100	100	100		
FOS/LAT	100	88.3	71.6	N.T	89.0	N.T	N.T	N.T	N.T	N.T	N.T	N.T	N.T	N.T	N.T	N.T	N.T	N.T	N.T		
FOS/NaCl	100	89.7	85.6	71.7	97.2	95.2	94.2	94.1	N.T	100	100	98.5	100	100	100	100	100	100	100		

W = week

Filling volume = 0.5 ml

FOS = Phosphate buffer

SAC = Sucrose, GLY = glycine, GLU = glucose, MAN = mannitol, LAT = lactose

N.T. = not tested

TABLE 9 - COMPATIBILITY WITH DIFFERENT EXCIPIENTS

Bioassay data (IU/ml)

LOT	T=0	50° C				40° C				25° C		4° C	
		1 W	2 W	7 W	2 W	4 W	7 W	10 W	8 W	11 W	8 W	12 W	8 W
FOS/SAC	9473	7854	8245*	N.V.	8098	10368	9126	8269	8809*	11222	9588	8489	9588
FOS/GLY	7850*	5642	4913	-	6421	8112	6780	6635*	-	7159	-	6821*	-
FOS/GLU	8370	N.V.	-	-	-	-	-	-	-	-	-	-	-
FOS/MAN	9498	7031	7224	6321*	10605	13216	9374*	6904	7285	7941	10079	8762	10079
FOS/LAT	7976	N.V.	-	-	-	-	-	-	-	-	-	-	-
FOS/NaCl	8486	8394	6433	-	9262	10576		7578	9151*	9353*	8804	8377	8804

W= week

Filling volume=0.5 ml

* = one valid assay

n.v. = not valid assay

- = not tested

FOS: Phosphate buffer, SAC = Sucrose, GLY = glycine, GLU = glucose, MAN = mannitol, LAT = lactose

TABLE 10 - LIQUID FORMULATION: CONC. 5,000 IU/vial**HPSEC Stability data: purity (%)****Formulation development**

LOT	50°C			40°C
	T=0	1 W	3 W	3 W
HCG/5000/S01	100	90.0	86.3	97.2
HCG/5000/M01	100	89.5	88.3	97.6

W=week

S01=sucrose

M01=mannitol

TABLE 11 - LIQUID FORMULATION: CONC. 10,000 IU/vial**HP-SEC Stability data: purity (%)****Formulation development**

LOT	50°C			40 °C
	T=0	1 W	3 W	3 W
HCG/10000/S01	100	91.8	88.9	97.9
HCG/10000/M01	100	93.4	92.1	97.2

W=week

S01=sucrose

M01=mannitol

TABLE 12 - LIQUID FORMULATION **α subunit purity by RP-HPLC**

LOT		50°C
	T=0	1 W
HCG/5000/S01 α(%)	100	90.2
HCG/5000/M01 α(%)	100	94.7

W= week

S01 = Sucrose

M01= Mannitol

TABLE 13 - LIQUID FORMULATION **α subunit parity by RP-HPLC**

LOT		50°C
	T=0	1 W
HCG/10000/S01 α(%)	100	92.4
HCG/10000/M01 α(%)	100	95.1

W= week

S01 = Sucrose

M01= Mannitol

TABLE 14 - LIQUID FORMULATION
Bioassay data (IU/ml)

LOT	30°C				
	T=0	1 W	3 W	4 W	5 W
HCG/5000/S01	09194 (7484-11298)	6427 (4770-8660)	6757 (5454-8371)	-	N V
HCG/5000/M01	8548 (6376-11459)	9249 (7495-11414)	6977 (5649-8618)	6207 (4767-8082)	3219* (1436-5150)

LOT	40°C			
	4 W	6 W	10 W	13 W
HCG/5000/S01	8682* (6082-12393)	10102 (7733-13195)	8192 (6276-10692)	-
HCG/5000/M01	10203 (7813-13325)	7959 (6118-10356)	-	7309 (5932-9005)

W = Week

N V = not valid assay

S01 = Sucrose

M01 = Mannitol

* = one valid assay

TABLE 14 (CONT.)

LOT	T=0	25°C		
		5 W	13 W	24 W
HCG/5000/S01	9194 (7484-11298)	-	-	-
HCG/5000/M01	8548 (6376-11459)	6660* (3855-10118)	8969 (7007-11479)	8232* (5787-11712)

LOT	5 W	4°C		
		13 W	24 W	
HCG/5000/S01	7555 (5904-9667)	-	-	
HCG/5000/M01	8869* (5968-12826)	10330 (8167-13065)	9799 (7714-12447)	

W=week

NV= not valid assay

*=one valid assay

S01=sucrose

M01=mannitol

TABLE 15 - LIQUID FORMULATION : 10,000 IU/VIAL

Bioassay data (IU/ml)

LOT	50 °C			
	T = 0	1 W	2 W	4 W
HCG/10000/S01	20273 (15170-27091)	15531 (11842-20368)	14971 (11307-19824)	-
HCG/10000/M01	18919 (14150-25295)	15880 (12605-20006)	13495 (9994-18222)	14855 (11579-19058)

LOT	40 °C		
	4 W	6 W	13 W
HCG/10000/S01	22201 (16648-29607)	14977 (12075-18576)	-
HCG/10000/M01	19508 (14201-26797)	14680 (11328-19022)	14606 (11580-18423)

W = Week

N V = not valid assay

S01 = Sucrose

M01 = Mannitol

* = one valid assay

TABLE 15 (CONT.)

LOT	25°C				
	T = 0	5 W	10 W	13 W	24 W
HCG/10000/S01	20273 (15170-27091)	-	17812* (11809-26112)	-	-
HCG/10000/M01	18919 (14150-25295)	17890 (14467-22122)	15494 (12638-18996)	16419 (12890-20915)	18991 (15311-23556)

LOT	4°C		
	5 W	13 W	24 W
HCG/10000/S01	21616 (17596-26555)	-	-
HCG/10000/M01	20666 (17390-24559)	17096 (13503-21646)	19553 (14494-26377)

W = Week

N V = not valid assay

S01 = Sucrose

M01 = Mannitol

* = one valid assay

CLAIMS

1. A liquid pharmaceutical composition comprising human Chorionic Gonadotropin and a stabilising amount of a polyalcohol or a non-reducing sugar.
5
2. A liquid pharmaceutical composition according to Claim 1, wherein the polyalcohol is mannitol.
- 10 3. A liquid pharmaceutical composition according to Claim 1, wherein the non-reducing sugar is sucrose.
4. A liquid pharmaceutical composition according to Claim 1, wherein the human Chorionic Gonadotropin is recombinant.
15
5. A liquid pharmaceutical composition according to any of Claims 1 to 4, wherein the solution is a buffered aqueous solution.
6. A liquid pharmaceutical composition according to Claim 5, wherein
20 the buffer solution is selected from the group consisting of acetate, succinate and phosphate buffer.
7. A liquid pharmaceutical composition according to Claim 6, wherein the buffer is phosphate buffer.
25
8. A liquid pharmaceutical composition according to any of Claims 5 to 7, wherein the buffer solution is at pH 7.00.
9. A liquid pharmaceutical composition according to any of Claims 5 to
30 8, wherein the buffer solution is 0.01 M.
10. A liquid pharmaceutical composition according to Claim 1, comprising from 1,000 to 40,000 IU/ml of hCG and from 10 to 180 mg/l of mannitol in a 0.01 M phosphate buffer at pH 7.00.
35

11. A process for the preparation of a liquid pharmaceutical composition according to Claim 1, comprising diluting a hCG bulk solution in a buffer solution containing the excipients.
- 5 12. A form of presentation of a liquid pharmaceutical composition of Claim 1 hermetically closed in a sterile condition in a container suitable for the storage before the use.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 95/01055

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K47/26 A61K38/24

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,93 11788 (APPLIED RESEARCH SYSTEMS ARS HOLDING N.V.) 24 June 1993 cited in the application see page 2, line 27 - page 3, line 32	1-9,11
Y	see page 7, line 8 - line 19; claims 1-14 ---	10,12
X	EP,A,0 448 146 (AKZO N.V.) 25 September 1991 see page 2, line 28 - line 45 see page 4, line 15 - line 30 see page 5, line 10 - line 12 see page 5, line 50 ---	1,3,5-8
Y	WO,A,89 04177 (GENENTECH, INC.) 18 May 1989 see page 2, line 32 - page 3, line 10 see page 4, paragraph 2 ---	10,12
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"A" document member of the same patent family

Date of the actual completion of the international search

29 June 1995

Date of mailing of the international search report

04.08.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Tzschoppe, D

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 95/01055

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP,A,0 597 101 (HAGIWARA, HIDEAKI) 18 May 1994 see abstract -----	10, 12

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No
PCT/EP 95/01055

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9311788	24-06-93	IT-B- 1250075	30-03-95
		AU-B- 3266293	19-07-93
		CA-A- 2124801	24-06-93
		EP-A- 0618808	12-10-94
EP-A-448146	25-09-91	AT-T- 107172	15-07-94
		AU-B- 631730	03-12-92
		AU-A- 7362391	26-09-91
		DE-D- 69102465	21-07-94
		DE-T- 69102465	02-03-95
		ES-T- 2057731	16-10-94
		JP-A- 4217630	07-08-92
		US-A- 5384132	24-01-95
		US-A- 5270057	14-12-93
WO-A-8904177	18-05-89	AU-A- 2724588	01-06-89
		CA-A- 1335176	11-04-95
		DE-D- 3888197	07-04-94
		DE-T- 3888197	18-08-94
		EP-A- 0386106	12-09-90
		IE-B- 60875	24-08-94
		JP-T- 3500882	28-02-91
		US-A- 5151265	29-09-92
EP-A-597101	18-05-94	JP-A- 5025058	02-02-93
		AU-A- 2330592	23-02-93
		WO-A- 9301835	04-02-93